

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 November 2004 (11.11.2004)


PCT

(10) International Publication Number
WO 2004/097024 A1

- (51) International Patent Classification⁷: C12N 15/82, C07K 14/415
- (21) International Application Number: PCT/SE2004/000604
- (22) International Filing Date: 20 April 2004 (20.04.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0301233-3 28 April 2003 (28.04.2003) SE
- (71) Applicant (for all designated States except US): SWE-TREE TECHNOLOGIES AB [SE/SE]; P O Box 7981, S-907 19 Umeå (SE).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): HERTZBERG, Magnus [SE/SE]; Istidskatan 27, S-906 55 Umeå (SE).
- (74) Agent: BERGENSTRÄHLE & LINDVALL AB; P.O. Box 17704, S-118 93 Stockholm (SE).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,

[Continued on next page]

(54) Title: TISSUE SPECIFIC PROMOTERS



(57) Abstract: Tissue specific promoters (SEQ. ID. NO. 1 - 5) are disclosed, these being preferentially expressed in the xylem tissue of woody plants. These promoters are used e.g. to specifically regulate gene expression in xylem tissue and alter xylem properties in plants. Transgenic plants, exhibiting modified xylem properties can be produced using these promoters.

WO 2004/097024 A1



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *with international search report*

Tissue specific promoters

The present invention relates to the field of gene technology, and in particular promoters driving gene expression during the development of xylem in any plant, and
5 in particular in the cambial region of a woody plant. The invention makes available such promoters and their practical use.

Background of the invention

Xylem formation in plants is an important process developmentally, and it is also very important in the production of fibres that are of economic value. Wood formation in
10 trees is an economically extremely important example of this process. It is the result of a series of defined developmental events from the initial cell division, through cell expansion and secondary wall formation ending in a zone of apoptosis. Wood is produced from the vascular cambium which is a lateral meristem undergoing terminal differentiation ending in autolysis of the cell protoplast.

15 The present inventors have in the studies leading to the present invention used Poplar that is a well-established model for woody perennials. It has a small genome, only about five times larger than Arabidopsis, another frequently used model. Further, Poplar is a species that is used extensively for forward genetics. Poplar and Arabidopsis are also phylogenetically closely related and access to over 100 000
20 Poplar ESTs combined with the complete sequence of the Arabidopsis genome make the combination of these two models an excellent choice for studies of xylem differentiation.

When and where a gene is active is important for its function. One level of regulation is the control of transcriptional initiation. The transcriptional expression of a gene is
25 controlled by the cis-acting DNA elements located mainly (but not exclusively) upstream (5') of the coding sequence. When using gene technology and GMOs (Genetically Modified Organisms) as tools in breeding and product development, there are two basic parts: the gene which gives the wanted effect; and the cis-acting DNA regulatory sequence which determines when, where and the level of gene
30 product expressed. Regulation also occurs at levels such as transcript stability, translational -initiation and -progression, protein activity and so forth. There are genes that have promoters driving their expression during different and specific

stages of xylem formation. These promoters will be essential in specifically directing the effect on any gene when genetically modifying xylem properties and altering the amount of xylem produced. The number of available promoters is limited, which means that new ones will be of great importance when tuning and fine tuning the expression of genes in the study and manipulation of xylem formation.

Modification of specific processes during cell development in higher species is also of great commercial interest, not only when it comes to improving the properties of trees, but also in other plants whose fibres have commercial use. New tools, such as promoters for such modification, are needed.

10

Prior art

Fredrik Sterky *et al.* (Proc. Natl. Acad. Sci. USA, 1998 (95), 13330-13335) and Magnus Hertzberg *et al.* (Proc. Natl. Acad. Sci. USA 2001 (98) 14732-14737) have published the results of a large-scale gene discovery program in two poplar species, comprising 5,629 ESTs from the wood forming tissues of *Populus tremula* L. x *tremuloides* Michx. and *Populus trichocarpa* 'Trichobel.' These ESTs represented a total of 3,719 unique transcripts for the two cDNA libraries and putative functions could be assigned to 2,245 of these transcripts. The authors state that the EST data presented will be valuable in identifying genes involved in the formation of secondary xylem and phloem in plants, but fail to give clear directions as to how the identification could be performed.

20.

In the prior art (Hertzberg *et al.*) transcript profiling of cDNA at defined stages of xylogenesis were performed using cDNA microarray analysis of a poplar unigene set of 2995 ESTs.

However, in order to make possible the practical use of the information published by Sterky *et al.*, and Hertzberg *et al.*, a new approach is needed.

25

U.S. 2002138870 A1 relates to a method of simultaneous transformation of plants with multiple genes from the phenylpropanoid pathways including 4CL, Cald5H, AldOMT, SAD and CAD genes and combinations thereof to produce various lines of transgenic plants displaying altered agronomic traits. The agronomic traits of the plants are regulated by the orientation of the specific genes and the selected gene combinations, which are incorporated into the plant genome. U.S. 2002138870

30

mentions the desirability of using tissue specific promoters, e.g. such that would confine the expression of the transgenes in developing xylem.

In general, it still remains to identify new, useful promoters. One particular problem remaining is how to identify the potentially most important genes and their
5 corresponding promoters, and to relate these to specific developmental properties of the cell. Another problem is to clone all relevant cis-acting transcriptional control elements so that the cloned DNA fragment drives transcription in the wanted specific expression pattern.

A particular problem is to identify specific promoters, related to specific cell types,
10 developmental stages and/or functions in the plant that are not expressed in other plant tissues.

Summary of the invention

The present invention offers a solution to the above problem by making available isolated nucleic acid sequences, each comprising a promoter sequence specifically
15 expressed in the xylem forming tissue in a plant, said sequence being chosen among SEQ ID NOs 1 through 5; sequences functionally homologous to SEQ ID NOs 1 through 5; and sequences showing at least 90% homology to SEQ ID NOs 1 through 5.

The sequences are presented in the figures, and in the attached sequence listing,
20 prepared using the prescribed software (PatentIn 3.1).

Further, the present invention makes available transgenic plants, methods for their production, and seeds and seedlings of such plants, as well as a method for expressing specific genes in the xylem of a plant, and nucleic acid constructs, as defined in the attached claims, incorporated herein by reference.

25 Short description of the drawings

The invention will be illustrated in closer detail in the following description, examples and drawings, in which:

Figure 1 shows consensus sequences from the genome walk fragment. 1a) LMX2 A014P10U; 1b) LMX3 A044P26U; 1c) LMX4 A050P49U; 1d) LMX5 A055P19U; 1e)
30 LMP1 A001P79U, corresponding to the sequences SEQ ID NO 1 – 5.

Underlined sequence = promoter sequence (part of the GUS construct)

Sequence in italics = sequence from EST

ATG (bold) = Potential translational start codon

Figure 2 shows a cross section of a hybrid aspen stem, stained with Toulidine blue. The sample positions are indicated with black bars, below the cross section.

- 5 **Figure 3** shows the expression patterns for the 5 ESTs from which the promoters have been cloned. This corresponds to the main expression caused by the respective promoters.

Figure 4 shows the results from northern blot analysis of expression levels of the selected genes in different plant tissue:

- 10 Lane 1: Apical shoot: Top shoot tip of about 1 cm.
- Lane 2: Leaf veins: The main veins were collected from old leaves.
- Lane 3: Elongating stem: The internodes below the apical shoot, to the eight node of the stem. All leaves and buds were removed.
- Lane 4: Old leaves: Healthy leaves more than 15 cm in size. The petiole and
- 15 main veins were removed.
- Lane 5: Phloem: 15 cm stem pieces were collected. These were used for phloem and xylem preparation by scraping the xylem and the inner-side of the bark.
- Lane 6: Root: The 2 – 3 cm apical part of root tips without any root hairs.
- 20 Lane 7: Young leaves: Leaves less than 12 cm in size. The petiole and main veins were removed.
- Lane 8: Xylem: Tissue collected as for the stem pieces (See Lane 5), the wood side being scraped.

25

Description

Definitions

In the description, the term "xylem", as in xylem forming tissue, is intended to encompass all the vascular tissue through which most of the water and minerals of a plant are conducted. The principle conducting cells of the xylem are the tracheids and

the vessel elements, or vessel elements. Both are elongated cells that have secondary walls and lack protoplasts at maturity. Xylem tissue also contains parenchyma cells that store various substances.

5 The term "vascular cambium" is meant to encompass the vascular meristem mother cells, the division of which produces secondary xylem and secondary phloem.

The terms "wood" or "wood forming cells" are meant to encompass all cells whose metabolic events and processes are involved in the development of the vascular meristem and all of its end products. The term "wood" is also used in its most general sense, encompassing both softwood and hardwood.

10 The term "wood formation properties" encompasses both biological, chemical and physiological properties, for example but not limited to cell elongation, cell expansion, apoptosis, carbohydrate composition, fibre length, fibre thickness, and other fibre properties.

15 "Homology" is here used as a measure of the similarity between sequences; the greater the sequence homology between two sequences, the greater degree of hybridisation. Hybrid formation can be measured by a person skilled in the art, using known procedures.

In the present invention, sequences sharing at least 90 %, preferably at least 91%, more preferably at least 92%, more preferably at least 93%, more preferably at least 20 94%, more preferably at least 95%, more preferably at least 96%, more preferably at least 97%, more preferably at least 98%, and most preferably at least 99 % homology are encompassed.

25 "Functionally homologous" means sequences sharing perhaps a lower structural homology with the disclosed sequence, but exhibiting homologous function *in vivo*, in either the healthy or the diseased organism, e.g. coding the same or highly similar proteins with similar cellular functions. The present invention encompasses sequences functionally homologous to the sequences disclosed.

"Complementary" in the context of this description refers to the capacity for precise pairing between two nucleotides.

30 Further, in the context of the present invention, "hybridization" refers to hydrogen bonding, which may be Watson-Crick, Hoogsteen or reverse Hoogsteen hydrogen

bonding, between complementary nucleoside or nucleotide bases. Thus complementarity and hybridisation are terms used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target.

5 An antisense compound is specifically hybridisable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-specific target sequences under conditions in which specific binding is desired.

10 The phrase "hybridisation under stringent conditions" refers to criteria regarding temperature and buffers well known to those skilled in the art. See e.g. Sambrook, J., Fritsch, E.F. and Maniatis, T., Molecular cloning: A laboratory manual, 2nd Edition, Cold Spring Harbour Laboratory Press, USA (1989).

"Functionally inserted" or "operationally inserted" denotes that a sequence has been
15 inserted in a host genome in such orientation, location and with such promoters, where applicable, that the correct expression of said sequence occurs.

"Modulation" as used in this context means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, increase is the preferred form of modulation of gene expression and mRNA
20 is a preferred target.

Detailed description

The selection of developmental stage specific expressed genes was used to identify promoters that direct gene expression to specific cells within the xylem forming tissue. When cloned and analysed, these promoters were shown to be tissue
25 specific. Being in possession of these promoters, it becomes possible to drive the expression of any gene of interest in specific stages of xylem development and effect the production and properties of the xylem (wood). The methods to alter expression include ectopic and over-expression of the gene of interest, antisense regulation, RNA interference, gene silencing, the use of ribozymes etc.

30 Libraries were created from regions within the xylem forming tissue, representing the areas of cell division (A), cell expansion (ABC); secondary cell wall formation (CDE); cell apoptosis (E). See Fig. 2.

Tissue samples were prepared by taking tangential sections from the cambial region of the stem. Transverse hand-sections were collected in parallel to allow later identification of the exact location and tissue content of each sample. The individual sections measured 30 µm x 2 mm x 20 mm corresponding to a fresh weight of approximately 0.5 mg. Sections were pooled together into different developmental zones as indicated in Fig. 2. The selection of different zones was based on the radial diameter and anatomical features of the cells. Within these zones, the gene profiles showed a very high degree of tissue specificity.

The present invention discloses gene promoters and their use, said promoters identified via the characterisation of cDNA microarray analysis, selected from a high number of sequences, found in undifferentiated meristematic cells in the cambium through to matured xylem elements ending in cells entering programmed cell death or apoptosis.

The present invention makes available isolated nucleic acid sequences comprising a promoter sequence specifically expressed in the xylem forming tissue in a plant, wherein said sequence is chosen among: SEQ ID NOs 1 through 5; sequences comprising one of said sequences SEQ ID NOs 1 through 5; sequences being functionally homologous to SEQ ID NOs 1 through 5; or sequences showing at least 90% homology to one of SEQ ID NOs 1 through 5, preferably at least 91%, more preferably at least 92%, more preferably at least 93%, more preferably at least 94%, more preferably at least 95%, more preferably at least 96%, more preferably at least 97%, more preferably at least 98%, and most preferably at least 99 % homology.

The above definitions do not exclude each other, but may act together. For example a sequence exhibiting lower homology, e.g. 80 % homology or 70 % homology, but being functionally equivalent to these sequences, is included in this definition. The invention also encompasses sequences capable of hybridising under stringent conditions to at least one of these sequences.

One aspect of the invention is nucleic acid sequences as defined above, wherein the promoter sequence is expressed in a higher plant. Preferably said plant is a woody dicotyledon. Examples of plants include but are not limited to hardwood species, such as poplar, aspen, birch, willow, eucalyptus, sweetgum (liquidamber) etc., and softwood species (conifers) spruce, larch, hemlock and pine etc. Other plants of interest are so called fibrous plants, for example, but not limited to cotton, hemp,

sisal, flax, etc. The promoter could be used in other economically significant plants like wheat, maize, potatoes, oil seed rape etc in order to drive the expression of genes during vascular development.

5 The present invention makes available transgenic plants exhibiting modified properties when compared to the wild-type of said plant, e.g. modified wood formation properties, modified apoptosis properties, altered rooting properties, altered flowering or leaf pattern etc in comparison to the wild-type of said plant, wherein at least one of the above sequences is functionally inserted into said transgenic plant.

10 The transgenic plant is preferably a woody plant, most preferably a woody dicotyledon. Examples of transgenic plants include but are not limited to hardwood species, such as poplar, aspen, birch, willow, eucalyptus, sweetgum (liquidamber) etc., and softwood species (conifers) spruce, larch, hemlock and pine etc. Other plants of interest are so called fibrous plants, for example, but not limited to cotton, hemp, sisal, flax, etc. The promoter could be used in other economically significant
15 plants like wheat, maize, potatoes, oil seed rape etc in order to drive the expression of genes during vascular development.

The present invention also makes available a method for expressing specific genes in a specific zone of the plant, preferably the xylem of a plant, wherein at least one of the above sequences is used, functionally inserted into the plant. Consequently, the
20 present invention makes it possible to regulate the xylem forming tissue development process in a plant, and in particular woody plants, wherein a promoter as defined above is used.

The invention also encompasses plant reproduction products of the plants of the invention, carrying the sequences in their genome, e.g. seeds, fruits, cuttings and
25 parts of the plants, such as protoplasts, plant cells, calli or roots.

An intermediate, also encompassed by the invention, is a nucleic acid construct comprising a sequence as defined above. Such construct preferably comprises a vector chosen among a plasmid, a cosmid, a virus or a bacteriophage.

30 The present invention also includes a method for the production of a transgenic plant exhibiting modified wood formation and/ or apoptosis properties compared to the wild type of said plant, wherein a promoter as defined above is functionally inserted in the plant.

Those skilled in the art are well able to construct vectors and design protocols for recombinant gene expression.

Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, translational leader sequences, terminator fragments, polyadenylation sequences, enhancer sequences, effector genes, marker genes and other sequences as appropriate. For further details see, for example, Molecular Cloning: a Laboratory Manual: 2nd edition, Sambrook *et al.*, 1989, Cold Spring Harbor Laboratory Press.

Many known techniques and protocols for manipulation of nucleic acids, for example in preparation of nucleic acid constructs, mutagenesis, sequencing, introduction of DNA into cells and gene expression, and analysis of proteins, are described in detail in Ausabel *et al.* (Eds.), Current Protocols in Molecular Biology, 2nd edition, John Wiley & Sons, 1992.

Selectable genetic markers may be used consisting of chimaeric genes that confer selectable phenotypes such as resistance to antibiotics. Such selectable markers can include kanamycin, hygromycin, phosphinothricin, chlorsulfuron, methotrexate, gentamycin, spectinomycin, imidazolinones, d-aminoacids and glyphosate.

Nucleic acid molecules and vectors according to the present invention may be provided isolated and/or purified from their natural environment, in substantially pure or homogeneous form, or free of nucleic acid or genes of the species of interest or origin other than the sequence encoding promoter with the required function. Nucleic acid according to the present invention may include genomic DNA and may be wholly or partially synthetic. The term "isolate" encompasses all these possibilities.

Plants transformed with these DNA segments containing at least one of the sequences SEQ ID NO 1 – 5, a functionally homologous sequence to one of the said sequences, or a sequence exhibiting a homology of at least 90 % to one of said sequences, may be produced by standard techniques which are already known for the genetic manipulation of plants. DNA can be transformed into plant cells using any suitable technology, such as a disarmed Ti-plasmid vector carried by *Agrobacterium* exploiting its natural gene transfer ability, particle or microprojectile bombardment, microinjection, electroporation, other forms of direct DNA uptake, liposome mediated

DNA uptake. Physical methods for transformation of plant cells are reviewed in Oard, 1991, Biotech. Adv. 9: 1-11.

Following transformation, a plant may be regenerated, e.g. from single cells, callus tissue, leaf discs, as is standard in the art. Almost any plant can be entirely
5 regenerated from cells, tissues and organs of the plant. Available techniques are reviewed in Vasil *et al.*, Cell Culture and Somatic Cell Genetics of Plants, Vol I, II, and III, Laboratory Procedures and Their Applications, Academic Press, 1984; Weissbach, Methods for Plant Molecular Biology, Academic Press, 1989; or Clapham
10 *et al.*, Gene Transfer by particle bombardment and Embryonic cultures of *Picea abies* and the production of transgenic plantlets. Scandinavian Journal of Forest research 15 (2000) 151-160.

The particular choice of a transformation technology will be determined by its efficiency to transform certain plant species as well as the experience and preference of the person practicing the invention with a particular methodology of choice. It will
15 be apparent to the skilled person that the particular choice of a transformation system to introduce nucleic acid into the plant cells is not essential to or a limitation of the invention, nor is the choice of technique for plant regeneration, or the technique chosen for the later multiplication of the transformed plants.

Examples

20 1. The experimental system

Xylem development is a specialised form of plant development that has been studied in several systems including *Arabidopsis* and *Zinnia*. However, for *in vivo* studies the Poplar system is superior because of the highly organised structure of the Poplar stem allowing precise and accurate sampling of specific tissues at defined
25 developmental stages.

Secondary xylem is highly organised, and the development of different cell lines are strictly co-ordinated in space. The present inventors have separated different tissues at defined developmental stages from the cambial region using a cryotom, producing by tangential cryosectioning 30µm sections, with a fresh weight of about 0.5 mg. This
30 dissection technique gives samples highly enriched for cells at specific stages of differentiation as shown in Fig. 2. Five samples were collected to encompass the

ontogeny of wood formation: A) meristematic cells, B) early expansion, C) late expansion, D) early secondary wall formation and E) late stage of cell maturation.

Like other meristems, the main function of the vascular cambium (Fig. 2, zone A) is cell division and setting out patterns for differentiation. But unlike apical meristems, the vascular cambium contains two morphologically distinct initials; axially elongated fusiform initials and isodiametrical ray initials. Both types of initials divide periclinally to give rise to radial cell lines of phloem and xylem elements. Derivatives from the fusiform initials make up the axial cell system. In the wood of Angiosperm trees such as poplar this consists of two major cell types, vessel elements that have a function in water transport and fibres that give mechanical support to the stem. The ray initials give rise to the horizontally oriented ray cells that have a major function in lateral transport of carbohydrates supplied by the phloem sap and minerals supplied in the xylem sap. As the stem increases in diameter, more initials are required. Fusiform initials are formed by pseudotransverse anticlinal divisions, which result in a shortening of the daughter cells. Therefore fusiform initials both divide and elongate in the cambial meristem.

In the cell expansion zone, cambial derivatives enlarge with different polarities depending on cell type (Figure 2, zones B and C). Developing fibres expand radially and increase in length by tip growth, whereas vessels expand only in the radial direction. Vessel expansion is rapid and much more extensive than fibre expansion, giving these cells their morphological characteristics. Vessel differentiation also progresses at a much faster rate than the development of fibres and rays, which are synchronised. Vessels have thereby formed secondary walls already in zone C and completed their maturation in zone E. During the stages of division and expansion (zones A, B and C) there is a requirement for a continuous biosynthesis of primary walls, which in poplar mainly consist of cellulose, hemicellulose and pectin. Fibre elements and ray cells in contact with vessels (contact rays) differentiate slightly faster than fibres and ray cells without contact to vessels (isolation rays). Therefore, zone C contains a mixture of primary and primary + secondary walled cell types.

As soon as cell expansion is complete, the secondary wall is deposited in all xylem cells (zone D). One of the characteristic features of this stage of development is that the random organisation of cellulose microfibrils present in the primary wall now changes to a highly organised helical structure in the secondary cell wall. The

cellulose and hemicellulose framework of secondary walls is accumulating in several wall layers (S1, S2, S3) that can be recognised from the degree of cellulose microfibril orientation. At the final stage of secondary wall thickening the interspace within the cellulose and hemicellulose network is lignified, starting from the cell corners and progressing inwards. During the formation of the secondary wall, the xylem elements are also extensively sculptured to form the network of pits and pores, which allow for vessel/vessel and vessel/ray contacts. This activity involves patterned degradation of the primary wall. At the end of the maturation process (zone E), fibres enter programmed cell death, (vessels enters into programmed cell death earlier in zone C and D). In Contrast, ray cells continue to contribute to the lignification of the surrounding cells and remain alive in the sapwood for several years.

2. Promoter cloning and Sequencing

The present inventors have approached the problem of identifying tissue specific gene promoters by performing tissue specific transcript profiling. With cryo-sectioning it becomes possible to obtain samples that are highly enriched with specific types of cells. The organised nature of the tissue in connection with the cryo-sectioning technique provides samples that are highly enriched in specific cell types and thus enabled us to perform tissue specific transcript profiling.

The present inventors cloned the 5' upstream sequences of 5 genes in Hybrid aspen (*Populus tremula x tremuloides*). These genes are all expressed during specific phases of xylem (wood) formation as based on the cDNA microarray analysis (Hertzberg et al., 2001 PNAS). A number of selection criterias were set up to ensure the wanted result. The selection criteria included: the EST had to have a BLASTX hit in the Arabidopsis proteome indicating that the EST contained the full open reading frame (ORF), and thereby probably contains the full transcript; the EST had to have an expression pattern, based on cDNA microarray data (Hertzberg et al) that showed a clear differential expression in the regions of interest, and also had an strong hybridization signal suggesting that the promoter driving the expression of that EST is strong.

The five genes I - V are represented by the EST's given in Table 1 below. In the table, the name used for the cloned promoter fragment is also indicated, as well as the zone where the promoter is predominantly active.

Table 1.

GeneBank EST nr	Internal EST nr:	Expressed in zon:	Name of cloned promoter	Gene number in Fig. 1
AI162215	A014P10U	C-D	LMX 2	1 (Fig. 1a)
AI163548	A044P26U	C-D	LMX 3	2 (Fig. 1b)
AI163880	A050P49U	D-E	LMX 4	3 (Fig. 1c)
AI164126	A055P19U	C-D	LMX 5	4 (Fig. 1d)
AI161513	A001P79U	A-B	LMP 1	5 (Fig. 1e)

The expression pattern for the AI163548 EST is based on the expression pattern of AI162928 (A027P19U) which is an EST originating from the same gene.

- 5 The expression pattern for these genes in different parts of the plant were analysed with northern blot analysis. This data shows that the genes LMX2 through LMX5 all are predominantly expressed in secondary xylem. The LMP1 gene has a more general expression pattern but it has the highest signal in samples containing vascular cambium (3 and 5) as would be suggested from the microarray analysis.
- 10 These results indicate that these genes are predominantly expressed during cambial cell differentiation and not in other processes, which was a prerequisite for going further and cloning the 5' regions (promoters) of these genes.

Northern blot analysis, material and methods

- 15 Total RNA was prepared using a CTAB based method (Plant Molecular Biology Reporter volume 11 (2) 1993) and the RNA was subsequently cleaned by using the QIAGEN Rneasy® Plant Mini Kit. RNA concentrations were determined using Ribo Greer® RNA quantification kit (Molecular Probes Eugen, Oregon) and the quality of the RNA was checked by agarose gel analysis. Equal amounts of RNA (15 µg) from the different samples were separated on a formaldehyde agarose gel (Sambrook, J; 20 Fritsch, E, Maniatis, T., Molecular cloning: A laboratory manual, Cold Spring Harbour Press, New York, 1989). The RNA was then blotted to a Hybond-N+ filter according

to standard methods and UV-crosslinked to the filter. EcoR1/ Not1 fragments were recovered from the EST clones. The fragments were labelled using the Strip-EZ™ DNA labelling kit from Ambion. Hybridizations were performed in Church buffer at 65°C (Church, G.M. and Gilbert, W. Genomic Sequencing, Proc. Natl. Acad. Sci. USA 81. 1991-1995, 1984). The filters were then washed (at 65°C) twice for 15 min in 0.5 % BSA, 1 mM EDTA, 40 mM NaHPO₄ pH 7.2, 5 % SDS and four times for 5 min in 1 mM, EDTA, 40mM NaHPO₄ pH 7.2, 1 % SDS. The hybridization results were analysed using a phosphoimager (Molecular Imager System GS525). Each filter were used up to two times and stripped using the Strip-EZ™ DNA kit protocol before re-probing.

Promoter cloning and plant transformation

Upstream genomic DNA sequences were cloned using the Universal GenomWalker™ kit from Clontech according to the manufactures recommendations. 8 genome walker libraries were constructed and used. The longest or in some cases the most specific amplified fragment was cloned into pGEM-T-easy vector (Promega) and sequenced. After sequencing the EST clone and a presumed translational start codon (based on the start codon in the best blastx hit in the Arabidopsis proteome) were mapped on to the sequence (Fig. 1, see sequences 1 - 5). Based on this, the presumed promoter was cloned from hybrid aspen genomic DNA using PCR and subsequently ligated into pPCV812.km (R. Walden, C. Koncz, J. Schell. Methods in Plant Moll Cell Biol. 1, 175 (1990)) or into pGWFS7 (M. Karimi, D. Inze and A. Depicker. Trends in Plant Sciences 7 (5)193-195, 2002) producing a transcriptional promoter GUS reporter construct. The promoter fragment was designed from 10-50bp upstream the start codon to as far 5' as possible, giving 5' sequences ranging from 1093 to 1807 bp. These constructs were transformed into hybrid aspen clone T89 using agrobacterium and transgenic plants were generated (Nilsson O. et al., Spatial pattern of cauliflower mosaic virus 25S promoter luciferase expression in transgenic hybrid aspen trees monitored by enzymatic assay and non-destructive imaging. Transgenic research 1. 209-220 (1992)). Following root initiation the plants were potted in 50% expanded clay (Leca ~~linker~~, size 2-6 mm) and 50% soil (Weibulls kronmull, YrkesPlantJord) and grown in green house with 18h light and 6h dark periods using Osram Powerstar HQI BT 400W lamps as supplementary light.

GUS analysis

To visualise the expression of the GUS reporter gene, tissue section and parts were stained in X-gluc essentially according to Nilsson *et al* (The *Agrobacterium rhizogenes* rolB and rolC promoters, *Physiologia Plantarum* 100: 456-462 (1997)) and to Regan *et al.* (Accurate and high resolution *in situ* hybridization analysis of gene expression in secondary stem tissues, *Plant J.* 19, 363-369 (1999)). Initial results on the promoter LMX5 show vascular associated expression (both in Arabidopsis and in hybrid aspen) and results for LMP1 showed an vascular associated expression in Arabidopsis and an root- and apical-meristem associated expression in hybrid aspen during plant regeneration.

These results indicate that the cis acting elements necessary for xylem associated expression were included in these constructs, this is not obvious and is therefore a surprising result, when the localisation of these sequences could be both further upstream (5') and also 3' of the transcription start.

A further and more detailed analysis of the Lmp1 and Lmx5 promoters in larger 2-3 m high plants, performed during the priority year, revealed highly specific expression patterns for these two promoters respectively. In the stem of the Lmp1 Gus lines the Gus activity were mainly present in the cambium cells + the expanding developing xylem but also to a lesser extent in the young cells producing 2nd cell walls. Lmx 5 had the opposite expression pattern with the highest expression in young cells producing secondary cell walls and a lower expression in the cambium and expanding xylem cells. Both Lmp1 and Lmx5 are expressed in all the cells of the cambium and young secondary cell wall forming cells.

Both these promoters were active in the root tip and in the vasculature of leafs and roots. None of these promoters seem to have any expression in the apical meristem but they both have expression in young xylem, especially Lmp1 which is expressed close to the apical meristem in the developing vasculature. Lmx5 is also expressed in the developing vasculature but at an later stage.

Lmx2, Lmx3 and Lmx4 were analysed in Poplar transgenics in late stages of tissue culture. These three promoters were cloned into the pKGWFS7 vector. Lmx2 was expressed in vasculature of leafs stems and roots. It was also expressed in the root tip. In the stem it is seen that the main expression is in the cambium and all the way

into the 2nd cell wall forming cells. Lmx3 has an similar expression pattern with the notable difference that it is not expressed in the root tip but clearly in the vasculature of the root from about 0,5 to about 7 mm from the root tip. Lmx3 is also expressed in the apical meristem and young leaf primordia and are later in leaf and primary stem development confined to the vasculature. The Lmx4 gene was lowly expressed in the young tissues present in tissue culture, however an expression in the vasculature can clearly be seen, especially in the oldest vasculature present in the analysed samples.

Although the invention has been described with regard to its preferred embodiments, which constitute the best mode presently known to the inventors, it should be understood that various changes and modifications as would be obvious to one having the ordinary skill in this art may be made without departing from the scope of the invention which is set forth in the claims appended hereto.

References

- Ausabel *et al.* (Eds.), Current Protocols in Molecular Biology, 2nd edition, John Wiley & Sons, 1992
- Church, G.M. and Gilbert, W. Genomic Sequencing, Proc. Natl. Acad. Sci. USA 81.
5 1991-1995, 1984
- Clapham *et al* 2000, Gene Transfer by particle bombardment and Embryonic cultures of *Picea abies* and the production of transgenic plantlets. Scandinavian Journal of Forest research 15,151-160
- Hertzberg, M. *et al.*, Proc. Natl. Acad. Sci. USA 2001 (98) 14732-14737
- 10 Karimi, M., Inze D. and Depicker, A., Trends in Plant Sciences 7 (5)193-195, 2002
- Nilsson O. *et al.*, Spatial pattern of cauliflower mosaic virus 25S promoter luciferase expression in transgenic hybrid aspen trees monitored by enzymatic assay and non-destructive imaging. Transgenic research 1. 209-220 (1992)
- Nilsson *et al.*, The Agrobacterium rhizogenes rolB and rolC promoters, *Physiologia*
15 *Plantarum* 100: 456-462 (1997)
- Oard, 1991, Biotech. Adv. 9: 1-11
- Regan *et al.*, Accurate and high resolution *in situ* hybridization analysis of gene expression in secondary stem tissues, *Plant J.* 19, 363-369 (1999)
- Sambrook, J., Fritsch, E.F. and Maniatis, T., Molecular cloning: A laboratory manual,
20 2nd Edition, Cold Spring Harbour Laboratory Press, USA (1989)
- Sterky, F. *et al.*, Proc. Natl. Acad. Sci. USA, 1998 (95), 13330-13335
- Vasil *et al.*, Cell Culture and Somatic Cell Genetics of Plants, Vol I, II, and III, Laboratory Procedures and Their Applications, Academic Press, 1984
- Walden, R., Koncz, C., Schell, J., Methods in Plant Moll Cell Biol. 1, 175 (1990)
- 25 Weissbach, Methods for Plant Molecular Biology, Academic Press, 1989

Claims

1. An isolated nucleic acid sequence comprising a promoter sequence specifically expressed or active in the xylem forming tissue in a plant, **characterized** in that said sequence is chosen among:
 - 5 - SEQ ID NO 1 through 5
 - sequences being functionally homologous to any one of SEQ ID NO 1 through 5
 - sequences showing at least 90% homology to any one of SEQ ID NO 1 through 5.
- 10 2. The nucleic acid sequence according to claim 1, wherein said sequence is expressed or active in specific stages of xylem formation in a plant.
3. The nucleic acid sequence according to claim 1 or 2, wherein the promoter sequence is expressed or active in a plant.
4. The nucleic acid sequence according to claim 1 or 2, wherein the promoter
15 sequence is expressed or active in a woody plant or a fibrous plant.
5. The nucleic acid sequence according to claim 1 or 2, wherein the promoter sequence is expressed or active in a woody plant, said woody plant being a dicotyledon.
6. The nucleic acid sequence according to claim 1 or 2, wherein the promoter
20 sequence is expressed or active in a plant chosen among poplar, aspen, birch, willow, eucalyptus, sweetgum (liquidamber), spruce, larch, hemlock, pine, cotton, hemp, sisal, flax, wheat, maize, potatoes, and oil seed rape.
7. A transgenic plant exhibiting modified wood formation properties in comparison to the wild-type of said plant, **characterized** in that at least one of
25 the sequences according to claim 1 or 2 is functionally inserted into said transgenic plant.
8. A transgenic plant exhibiting modified apoptosis properties in comparison to the wild-type of said plant, **characterized** in that at least one of the sequences according to claim 1 or 2 is functionally inserted into said transgenic plant.

9. A transgenic plant according to claim 7 or 8, wherein said transgenic plant is a woody plant or a fibrous plant.
10. A transgenic plant according to claim 7 or 8, wherein said transgenic plant is chosen among poplar, aspen, birch, willow, eucalyptus, sweetgum
5 (liquidamber), spruce, larch, hemlock, pine, cotton, hemp, sisal, flax, wheat, maize, potatoes, and oil seed rape.
11. A method for expressing specific genes in the xylem of a plant, **characterised** in that at least one of the sequences according to claim 1 or 2 is used.
12. A method for production of a transgenic plant, **characterised** in that at least
10 one of the promoters according to claim 1 is functionally inserted into the plant.
13. Propagating material of a transgenic plant according to any one of claims 7 through 10, said propagating material carrying said at least one sequence in its genome.
14. Propagating material according to claim 13, said propagating material chosen
15 among seeds, fruits, cuttings and parts of the plants, such as protoplasts, plant cells, calli or roots.
15. A nucleic acid construct comprising a sequence according to claim 1 or 2.
16. A nucleic acid construct according to claim 15, wherein the construct comprises a vector chosen among a plasmid, a cosmid, a virus or a
20 bacteriophage.
17. A nucleic acid sequence capable of hybridising under stringent conditions to at least one of the sequences according to claim 1 or 2.
-

LMX2 A014P10U – SEQ ID NO. 1

>ATCCCTCCTATTAAATCCATCTAATTGTATGGAAAGGATGGATTTATAAAATTACATAGGTTTCAT
TGATAATATTTCCAATATTTAAAATTAATGGTCATCATCATTGATTAAATCACAATATTTAGTCCT
GTCTAATCATAGAAATCTTGTCTCCGTACATGAAAGTTGATTAATAAATAAGTTATTAATCTTAT
TAGTTAATTTGATGTATTGATTCATGAGTTTAGTAGATTTATCCATCAAATCTAATATGTATTTTTT
AGTTTTTGGTTCATTGATTTTTCTAATTGTTTTTTATTTTACTATTTTTCAGTATTGrATTGTTTTAAAA
AATAAAATTTATAGTTTATTTTTTTTTCTATAGAGCTATCACTATCTCAATAAATATCTTTTTAGAGT
TGGTGTITGATTTTGTAACATCAAATTTTATCATAAAATTAATAAAAAATAATTTATAAAAAACAACA
AAGTTTAAACCTATTGAAATCTATAACCCAAGTCACAAAGTTACCAAACTTATTTAATTTGGTATC
ATCTCAATAATAATTAAAAAATTATTGTTTTAATTTTTTTTTTAAATCTATCATATTTTAACTAATTATT
AAATTATCTTTGAACTTATTAAATTAATTTTCATGCTTGTTAGAGAAAATTTCTACTAATAAATAA
CTTTTGTCTACTAAGTCATGAAAAATGTGATCTTTAATGATACAAGAACAGAAGTCCAAAAGCAG
GGGGGGTAAAGAAAGGGAGAGGTTGCCTATATATTTGCCTTACACTTACTCTACAAGGAACCTTT
TGCACCTAATGAGCGTAATAAATAGTATAGCAyGATAGCAATAGCACAAATAGCTCAGAAATGATAT
AGCTTCTAACTAATATAGAAGACAACACGAGCCCAAGATGTTAGCCCTCCAACCTTATGCCACGCTA
ACCTGCGTTTATACTCTAAAAAGATAAATGTAGAAAAAAAGTGAAAACCTAACTGTACATGGCTTTA
GCTTACTCCTACTCAAACAGCCATAAAAAATCCCAAAACGAGGCCTCTAAAAGTGTCTCATCAT
TCTGTTGCTTCAGTTAAGCTTCTGTTCTCTGCCTCCAACAATGAGCTTAGCAGCCTCTCAGTCCGC
TGCAATGAAACCTTCGAAAGGGAACTTCCCATTTCGGTTGTCACTGTATAGTTTGCAGCTTTGC
ATTCAATTGCTCTCTTGATACGACACTCCAGGTCTGTTTTCAAGC

Fig. 1a

LMX3 A044P26U – SEQ ID NO. 2

>AAATTGCTTAATGGATAATATAATTAACCTAATTATTTATATACACCGGTCACCTTTCTAGTTTATG
TAACTACATATTTTGAATTTTCAATTCCTTTTCACTTTCTTTTGATGGAAAAGTTTTGAATTCATAATAA
TTGGATTGGACTTGTCTATTTTCTTTAGAGATTTATCAATTGTGTTTTATTTTTTAACATTTATTAAAT
TAAATGGACTAGGACTATACATACCTTATCTATGATGTATTTAGTATTGCGTTTCAAATAGGGGTTG
AATAAATTTTGAATTTTATTTCTGTTTAAAATTATTTTTTTATGTTTTCAAATCGTTTTGATGTGCTG
ATATCAAAAATAATTTTGAAAAAAATATATATTATTTGATGTATTTTGAATAAAAAATACCTTGAAA
AACAACTGGTTATTACAGTCTCAAACACTCCTAGTAGTCTTAGGCATAGTTTAAAACTCAGTCCGA
CCTGGCGGGTCGACTTGGGACTGGAACCGGGCCGAGTTAAAGAAAAAATAaAAAAAGAAAAAA
CTTGGTGTGACCTGGTGACTCGGTTGACCCAGATGCAACCCATTGATTTTTGTTTTTTTAAACTA
AAATTACATTGTTTTGCTTTTTTAAAAAATAATTGACCCGTCAAACCTGGTCAAAACCCGGAATC
GAGGTCTTGGACCGGGACGGGTCTAAAACTATGATCTTTGGTCAATATATTAAGACAACAAGGT
CAAATTAAGGGTAATCAAAGGAGAGTTTTTTTTGGTTGGTTGGTTGGTTTTGGACCTAATCATA
CAATCATAAAATAATCATGGATCAAATGTAGGTAATCAATCATGGTGTGTAATAGATTTTTTATA
TGATCATGTGTCACCAGCTAGGTCTTTTTCCAAAATTCTAACAAGAGGTTGGTATATGTAATAAAAT
TATCTTTAAACAGACACAACCCACACCCTCTATAATTAATCTTAACAACAAGTTAATATAAAACAAT
TTTTTTTTTTGTCTAAATATATTTGCTGAYGGGTAGCATCCCACCAACTATCTGGTGCAGATAGC
AACTCAAGTCCTTCCCAATAAACACATTTTTCCATGATTCTTTCACTTCAAGTGACAACATAACGA
AGCAAAGCCTCGTCAAATTAATAAAAAAAAAAAAAAAAAAGGGTTCCGAATAATCATAATGTGGCTTTTGA
GTTTGCCAGTACCTGTCATCGACCGTTAGCTTCAACCTCCGCAAAAAACATACAGCTTCGCTAT
AAAACAACCTCCTTGTTTATTAATTGTTGTCTTCCAGTCTCTCAATCTCTCTCAAAGAATCCCTTCT
TTTTCTCAAACCTCAGGGGTCAGTATCTGAGCCAACAACCTTGGTTTTTGATCACGTTTTGAGGG
GTTTAGCTTGATGGGCTCTGTGGAAAGATCAAAGAGGAGAGTTTCAAGTTATGGAAGAAAGCTATAG
TTTCAATTTGGTTTGTGTTTGTGTCATGG

Fig. 1b

LMX4 A050P49U – SEQ ID NO. 3

>ATCAGCGCATCAAAATTATCTAAAAACATTAAAAGTATATTAATTTCAAGAAAAAATAAAAAAT
TTAAAAAAATTTACAAATAAATTTAAAAATTCAAATACAAAACAGATTGATATAGAATTTACTATTAAT
TATACCTCTATAAGCAGAGACATAATATTCTCTTTTGAGAAACAAAATATTATTATAATTGATATTCC
AACAAAAAACAAATCAATCTCTTTATTTTAGAAAGTAAATAACTTATAATTGTTTATAATTGTTTG
TAGATGGACGTCATGTTTGTCTGGGGGTAGACGCCCTCTTGGTTAAAAATAAAAAATAAATTT
TCTAATCTTATCCAAAATTTGTAACCTCTGyCAACTGAAATATAGTATTCTTAGAATTGATATTTAA
ATGAAATCCAAATGCTTAGAATTTATTTTAGCCATGTTGATGAGAAAGTCAAGCACCCCATTAATAA
AAATAAAATAAAATAAAAAATTACAAGGTAATTATGCACCTAGAAATTTTATTTTTTAATATAATAA
TACCCCTTTCTGACTGGAAAAGGCTGCGAGGTTGTCATTACACAACAAATATCAGCTATTTTCAGG
ATCCATCGCAGTACTCGCAATCTTCACATTTAGCAAAATCAGAGCAGCGTATGCCCTGTTTTTTC
ACCTTTTTATCGCACTAATCCTAGAGACGAACAGTGTTTATTGTTTTCTCTTAATATTTGACCTTTT
CACTTATTyGAAAAGTCAAATGTTAATTTCCACCGAGCTTTATCCTGTAAATAGCGTGTTTAATGCC
AGCTGTAAATAAGATAGATGATCGATTTTGTATTTACATTCTCTTATCTTAGACGGAAAGTTATTAA
AAAAATAAAATAAAATAGAAACCTACCTAAGATGTGAACTCTTGGCGTCTCTAGCTGTTTCTGTCT
CTAGAACAACCTCTACGAGTATAGCCGAAGCATATCCTAGCAAGCTTGAACCATATCTACTACCTC
CCTCCCGCTATATATATATAAACCCTCTTCTCCTAGAAAAAATAACCCTCGATCTTCTTTT
TTGTGCTTATGTTGATTTTGTGTTGATTTTCATATATCATGAGCAATATACCAAGATCTCTTGCAGAC
TCTTCTTTCACCCCTTTCAAGCTTGCCATTTCTGCAGCTGACCCTTGGCCTTTCTCTCTGTT (The
predicted start codon lies 102bp upstream.)

Fig. 1c

LMX5 A055P19U – SEQ ID NO. 4

>TACTATAGGGCACGCGTGGTCGACGGCCCGGGCTGGTACTCGTCTTCGTGCACCAAGATGCAA
 AAGCTGGAATGAGAAATGCGCTAAATGCGAAAAGACAGAGAGAGCGAATAAATCGTGCAAAAAAG
 GAGTGGGGTGGGTAAACGGGTTGAGCTAGAAGAAGAAAAGGGACAAGTGCACCTTTAGGAGGGGG
 GCAACCAGAGCGTAGATGATAATGGTTCATGTGGAACAACACACATGAGCAGTTGGTGAGAACT
 TGAATGAACCCTAACAGCCCCAACCAAAACCCGGAGCCACCCTTACCGAACCACTTCTAAAAGT
 ACACCATGCCTTTTTCTTGAGCyTGGTTGCACAGGGTGCCrGGTGGGGTTGTTTCGTTTTGGGTAA
 TCATGCGATAGTTTAAATACCTTTGCGATAATCATATCAATGGyGACTTTTAAGCACATGTTAGGTG
 CTCGGTTCTTATCTAAACATGGACATGGCmACAAGAGTTAATGCTAAAATAATATACGTACATACC
 TGTGAATGAATCGTCGCTGTCTTCTGATTATGGCTTCAAATTAATATGCAGATAAAACAAGTGTCGA
 TTTTTCAGTGAAGATTTTATGAAAGTGCCCGTTCTCCTTAAGATTACCTGTGAATGAATCATCCCTG
 TCAGGCTGATCCTGGGTGNTTTTTCCCATGrTGTTCGGAAGATATAATTATATAAATGATGGAATT
 TACATGAAATAAGTTTCAGTACATTCTTAGCAGAAAAGCAATATCGACGAAGACAAATGATGCTGT
 TTAAGACAAACTGGGGTAATATCAATTTACTAGTAAGAGATTTGkCTGCTTTTCTTAATTCTCAAGA
 AACTTyCACTAAAATGCACAGCCATGTTAAACAATTTACGTTCAACTTAAAACTAAAAGTGCAGGA
 TGGGTAGCTATCCAAGAATGATTAATAAACGATTTTAAACAAGAACTACGTTTTTGTAACTTAAT
 TTTGATGGGCATGTAGTTAAACAAGTATTTGTCAATTGATCAATTCAAGAGCCATGTCTGCATCATAAT
 TGTGGGAGTGGAGGAGGCTTTTGTTGCTAGGGAAGGAATGCCTTCTTAGTTCATGGCTTTGGACT
 TCGGACAAGGAGCGCATAGAATGGGGTTACCATTTTTGAAAAAATTACATTTGAACCTCCAAC
 ATTATCATGTATGTTTAAATCTACAATCCTCGTCCGCTAGAAGAAGTTTGGGTCAAAGTCATCCGC
 TCAATGTAAACCATGGAGAGGCAGGGACTAATTGAAAATAGTATGTTAGTTGGAGGGTCTGCATG
 TATTATGTCCAAACATTCTCTTATTATTCCTGTATCATCTCTGAGAAATTCATCCGAAAATAATAAAA
 CAAATGGCCTTTTTTAAAAAAGAAGCTGATGCATAGGATACCAAAAGCGCCTTGTCCATTAGGA
 GCGTCAGACTTTGAAAATAAGACCAATAATTCCTGTAAGCTATCATCTCATCTTTTATTTTGTG
 AACTTGTAGACGTAGGCTTTAAGCGTTCCATGATGTTCAAGTCAATGTTGCTGTCTACTTGATTAT
 GGAATTTAATTCATTGCGCTCATAAGAAGATAAAAGGATTATGACGTTGAAGAACTCTGGTCACTC
 CTTACTTACGGTCACATAAAAAACGATGCATCTTTCCCCACCAACCATCTTCAAGTGAACCCACTTT
 CCCTTGCAATTAGGTAAGGAGTATGGGTAAAGTCATCTTCATGAAATTAGTCCCCTAGTGGAGCTAA
 TTCTACTCACTCCATATTTACTCATTCCACTATATAACGCCCTCAACGACCATCCTCAAAGCAACC
 CAAACACCTTCTTCTCTTCTCTTTCTCCCGGAAAATGGCCGGTTTGATTTCAGATCAGTTC
 CTTGTGCAATCCTGGTAGTCTTGTGCACGGTGGTGCCATTTTGGCTAAAGATTACACTGTAGGA
 GATACTTCAGGCTGGGCAATTGGTATGGATTATAGCACCTGGACTAGTGGCAAGACCTTTTCAGT
 TGGCGACAGCCTTGGTGAGTCCCTTTCTCTCAGTACACGTTACTGATCAAGATTAGTACTTGATT
 ACTAATCAGGTTTCTGTTTAAATATATAATCTAGTGTTAACTACGGAGGAGGCCACACGGTGGATG
 AAGTGAGAGCCAGTGACTACAGCACATGCACTACAGGCAATGCAATCACTTCAGATAGCAGTGGT
 GCTACCACAGTAGCCCTCAAGACTGCCGGAACCCATTATTTCAATTTGTGGTGTTCCTGGCCACTG
 TGGGAGTGGCATGAAGGTTTCAGTCACTGTTGCAGCAGCAGGATCGAGCACAAGTCCCTCCTCC
 TCAGGAACTCCATCTTCTGATGGCACTACCACTTCTCCGGCCGGTAGTAATGTCACCAATTACAA
 GCCTTCATCCAACAA

Fig. 1d

LMP1 A001P79U – SEQ ID NO. 5

>CCTATGGAGTGGTCTACTGTTCTATATGCAGGGCCACAAGGATCTGGAAAAATGGAAAACTTC
ATAAATCTAGCTTGGAATCACCTATCATGCTACTCACTGGTCACTAGAGTGCTGTATGTACTATGA
AGTTCAATCCAGATGGAAATGTCATTGCATCAGGATCTCATGACAAGGAAATTTTCTATGGTAAAT
TCATGGGGAATGCAAAAACCTTCATGGTTATGAGAGGGGCACAAGAATGCAGTTTGGATCTTCACT
GGACTACTGATGGGTCCCAGATAATATCAGCCAGTCTGACAAAATTGTCAGGGTAGGATGTCTGA
GGCTGGAAAACAGATAAAGAAATCGGCAGAACACTCCTCTTTTGTGAATTCATGCTATGCTGCCTT
TCTGCAGGGGGCCTCCACTTGTTGTAAGTGGATCTGATGATGGAACTGCAAACTTTGGGATATG
CGCCCAAAGGGTGCCATCCAAACATTTCCAGATAAATAAAATCACAGCTGTCAGTTTCTCTGATGC
ATCAAGATAAGATATTCACAGGTGyTGTAGACAATGATATTAAGGCAAGGGACATCCGCAAGGGT
GAAGTGACCGTGGCACTTGAGGGCCATCAGGATATGATTACAGGTATGCAGTTAAGTCCTGATG
GCTCATATCTGCTTATGAATGGCATGGACAACAAGCTCTGCATTTTCGCCCATTTGCACCACAAAA
TCGTTGTGTGAAGATTTTTGAAGGGCACCAGCACAACTTTGAAAAGAACTTGTGAAAAGATAGCT
GGTCACCAGATGGAAGCAAGGTTACTAGTGCAGATCGCATGGCTTATATATGAATCTGTCAATGA
GTGTGTCTTCCACCTACTGAACCTATCTTTGGATCATGCAGCAGCGACAAACAGATTTATCTTGGG
GAAATCTGATTACTATTTTGTGAGTTTGCTTAAACAGATTGTTAGTAACAGTGAGGATATTyCGGG
GCTTGCTTGTGAGCNCCCAACAGAAAAGATTAAACAGGTTGTAATGGATACTATGTTACATTTTAC
TTAACTTTCTCATATCCTGCTGACATGTTTGCAGTTCTTTTTGCTTGTGGCGATCCTTTCAACT
GAAGTATGTACGCATGAATCAATTAGCTCAAAATGTATGTTGAGCAATATGCCTGGTGAATGTTT
GTTAAGATACGGAAAATTAAGTTACTGAACTTGTACTCTTTGATTTTGAAACAAGTTCCGACTATG
TTCAATTGTCCTAGCACGGTGATATTGATACGTGCAATTTTCAACTATTTTGTCTTGTTTTCAACTA
TTTCTGTCTTGATTTTCAAAGTCTTTTTGAAAATAAAAAACCCAAAAATATGTTTTCAGTGTCTTGT
AGCTAAATCTGTTGTACAACCAAGCCTCCCCGACTTGCTTTGTGAAAAAGAAAGATGGGAGCTGG
GGGTTTTGTGTTGATTATACAAAACCTTACGAGATAACTGATATACAAGCAAAAACACCCTCATATT
ATTGAATCTGCGACTTCTACAGTTCATTTCTGAGGAAGAATTAGGTGACTACTTCTGGACTAGAT
ATTTCCGTGTTCAATGCGTCGCAACAATTGATTTGTTGGCTGGATTATGAACATGATAATCACACT
ATAGAAAGACGACGATTATATTTCAAGCTATGGACAATGATGTTACCTGCTCGAAGATCAATACGC
TTGAAGGCCAAGATATGATCAsCAGGTATGCAGGCAGTCCAGATGGCTTCTTGGGATATGAGCTT
CAAGATTGTTCTTGAAAAGAACCATAAATTGTAAATTCTCTGGGAATATTTTTTTGTAAATTAT
CAGCATTGCTATTTTTTTTTTCAAATTCATTTAAATATTGCTATTGGGTTAGTAGTTCAGCGCAATTA
ACTTAGCTGGCAGTCTGGCATTAGCTTCAAATCATATCACAACGTTAAAAGCAGCGTTGCAACTA
GTCAGCCCACTAGCATTTGCTGGTGACCAATCAGATCATAAAAAGGAAAAGTACCCGTTGGCAAAC
CCTCTTCAACGGTCATCTCCCTGCTTCATGATATATATGCTTAGATATCCCAGTATCCCCTTCAA
CAAATTATCTCAATCTTCAAAACCCAGTCCCAGTTACAGCAGATCATCAAAAATCTCAAAATGGGT
CTCATAAACAATCACAAATCCyGCCAGTGACTGAATCCCACCACTCACTCCACAAATCTTCGT
ATATGCAACTACATTCTCTTATGTGCAGCC

Fig. 1e






Fig. 2

**Fig. 3**



WO 2004/097024

PCT/SE2004/000604

52669.ST25
SEQUENCE LISTING

<110> SWETREE TECHNOLOGIES AB
HERTZBERG, Magnus

<120> Tissue specific promoters

<130> MH52669

<150> SE0301233-3

<151> 2003-04-28

<160> 5

<170> PatentIn version 3.1

<210> 1

<211> 1093

<212> DNA

<213> Populus tremula x tremuloides

<400> 1
atccctccta ttaaattccat ctaattgtat ggaaaggatg gatttataaa attacatagg 60
tttrcattga taatatttcc aatattttaa attaaatggt catcatcatt cgattaattc 120
acaatattta gtcctgtcta atcatagaaa tcttgtctcc gtacatgaaa gttgattaaa 180
aataaataag ttattaatct tattagttaa ttgatgtat tgattcatga gtttagtaga 240
tttatccatc aaatctaata tgtatttttt tagtttttgg ttcattgatt tttctaattg 300
tttttttatt ttactatttt tcagtattgr attgttttaa aaataaaatt tatagtttat 360
tttttttttc tatagagcta tcactatctc aataaatatc ttttttagagt tgggtgtttga 420
ttttgtaaac atcaaatttt atcataaaat taaataaaaa taatttataa aaacaacaaa 480
gtttaaacct attgaaatct ataaccaag tcacaaagtt accaaaactt atttaatttg 540
gtatcatctc aataataatt aaaaaattat tgttttaatt tttttttaa tctatcatat 600
tttaactaat tattttaaatt atctttgaac ttattaaatt aaattttca tgcttgtag 660
agaaaatatt ctactaataa ataacttttg tcatactaag tcatgaaaaa tgtgatcttt 720
aatgatacaa gaacagaagt ccaaagcag ggggggtaaa gaaagggaga gggtgcctat 780

Sida 1

52669.ST25

atatttgcct tacacttact ctacaaggaa ccttttgcac ctaatgagcg taataaatag	840
tatagcayga tagcaatagc acaatagctc agaaatgata tagcttctaa ctaatataga	900
agacaacacg agcccaagat gttagccctc caacttatgc caccctaacc tgcgtttata	960
ctctaaaaag ataatgtag aaaaaaagtg aaaacctaac tgtacatggc tttagcttta	1020
ctcctactca aacagccata aaaaatccca aaacgaggcc tctaaaagtg tcctcatcat	1080
tctgttgctt cag	1093

<210> 2

<211> 1245

<212> DNA

<213> Populus tremula x tremuloides

<400> 2

gaattcataa taattggatt ggacttgtct attttcttta gagatttatc aattgtgttt	60
tatttttttaa catattattaa attaaatgga ctaggactat acatacctta tctatgatgt	120
atttagtatt gcgtttcaaa taggggttga ataaattttg aattttattt ctgtttaaaa	180
ttattttttt tatgttttca aatcgttttg atgtgtgat atcaaaaata attttgaaaa	240
aaatatatat tattttgatg tatttttgaa taaaaaatat tttgaaaaac aactggttat	300
tacagtctca aacactccta gtagtcttag gcatagttta aaactcagtc cgacctggcg	360
ggtcgacttg ggactggaac cgggccgagt taaagaaaaa aataaaaaaa gaaaaaactt	420
ggtgtgacct ggtgactcgg ttgaccaga tgcaacccat tgatttttgt tttttttaa	480
ctaaaattac attgttttgc ttttttaaaa aaaaaaattg acccgtcaaa cctgggtcaaa	540
acccggaatc gaggtcttgg accgggacgg gtctaaaaac tatgatcttt ggtcaatata	600
ttaagacaac aaggtcaaata taaagggtta tcaaaggaga gttttttttt ggttggttgg	660
ttggttttgg acctaatcat acaatcataa aataatcatg gatcaaatg taggtaattc	720
aatcatggtg tgtaatagat tttttatatg atcatgtgtc accagctagg tctttttcca	780
aaattctaac aagaggttgg tatatgtaat aaaattatct ttaaacagac acaaccaca	840
ccctctataa ttaatcttaa caacaagtta atataaaaca attttttttt tttgtctaaa	900
tatattttgc tgaygggtag catcccacca actatctggg gcagatagca actcaagtcc	960
ttccaataa acacattttt ccatgattcc tttcacttca agtgacaact aaacgaagca	1020
aagcctcgtc aaattaaaaa aaaaaaaaaa aggggtccga ataataataa tgtggctttt	1080
gagtttgcca gtacctgtca tcgaccrgtt agcttcaacc tccgcaaaaa aacatacagc	1140
ttcgctataa aacaactcct tgtttattaa ttgtgtctt ccactctctt caatctctct	1200
caaagaatcc ctttcttttt cctcaaaact caggggtcag tatct	1245

52669.ST25

<210> 3

<211> 1133

<212> DNA

<213> Populus tremula x tremuloides

<400> 3

```

atcagcgcac caaaattatc taaaaacatt aaaagtatat taatttcaag aaaaaaata 60
aaaaaattta aaaaaaattt acaaataaat ttaaaattca aatacaaac agattgatat 120
agaatttact attaattata cctctataag cagagacata atattctctt ttgagaaaca 180
aaatattatt ataattgata ttccaacaaa aaaacaaatc aatctcttta ttttagaaag 240
taaaataact tataattggt cataattggt tgtagatgga cgtcatgttt gtcgggggta 300
gacgccctct tgggttaaaat aaaataaaaa aataattttc taatcttatc caaaatttgt 360
aacctctgyc aactgaaata tagtattctt agaattgata ttttaaatga aatccaaatg 420
cttagaattt attttagcca tgttgatgag aaagtcaagc accccattaa ataaaataaa 480
ataaaataaa aaattacaag gtaattatgc acctagaaat ttttattttt ttaatatat 540
aataccctt tctgactgga aaaggctgag aggttgatc tccacacaaca aatatcagct 600
atttcaggat ccacgcagc actcgcaatc ttcacattta gcaaaatcag agcagcgtat 660
gccctgtttt tttcacctt ttatcgact aatcctagag acgaacagt tttattgttt 720
ttctcttaat atttgacct ttcacttatt ygaagagtca aatgttaatt tccaccgagc 780
tttatcctgt aaatagcgtg tttaatgcca gctgtaaata agatagatga tcgattttgt 840
atttacattc tcttatctta gacggaaagt tattaataaa ataaaataaa atagaaacct 900
acctaagatg tgaactcttg gcgtctctag ctgtttctgt ctctagaaca acctctacga 960
gtatagccga agcatatcct agcaagcttg aaccatatct actacctccc tccccgctat 1020
atatatataa accactcttc ttcctagaaa aaaaataacc ctcgatcttt ctttcttgtg 1080
cttatgttga ttttgtttga ttttcatata tcatgagcaa tataccaaga tct 1133

```

<210> 4

<211> 1807

<212> DNA

<213> Populus tremula x tremuloides

<220>

<221> misc_feature

<222> (616)..(616)

<223> unknown amino

52669.ST25

<400> 4
gatgcaaaaag ctggaatgag aatgcgctaa atgcgaaaag acagagagag cgaataaatc 60
gtgcaaaaaa aggagtgggg tgggtaacgg gttgagctag aagaagaaaa gggacaagtg 120
cacttttagga ggggggcaac cagagcgtag atgataatgg ttcatgtgga aacaacacac 180
atgagcagtt ggtgagaact tgaatgaacc ctaacagccc aaccaaaccg ggagccaccc 240
ttaccgaacc accacttcta aaagtacacc atgccttttt cttgagcytg gttgcacagg 300
gtgccrggtg gggttgtttc gttttgggta atcatgcat agtttaaata cctttgcat 360
aatcatatca atggygactt ttaagcacat gttaggtgct cggttcttat ctaaactggt 420
acatggcmac aagagttaat gctaaaataa tatacgtaca tacctgtgaa tgaatcgctg 480
ctgtcttctg attatggctt caaattaata tgcagataaa caagtgtcga tttttcagtg 540
aagattttat gaaagtgtcc gttctcctta agattacctg tgaatgaatc atccctgtca 600
ggctgatcct ggggtgnttt tccccatggt gttcggaaga tataattata taaatgatgg 660
aatttacatg aaataagttt cagtacattc ttagcagaaa agcaatatcg acgaagacaa 720
atgatgctgt ttaagacaaa ctggggtaat atcaatttac tagtaagaga tttgkctgct 780
tttcttaatt ctcaagaaac ttycactaaa atgcacagcc atgttaaaca atttacgttc 840
aacttaaaaa ctaaaactgc aggatgggta gctatccaag aatgattaat aaacgatttt 900
aacaagaac tacgtttttt gtaaaactta ttttgatggg catgtagtta acaagtattt 960
gtcattgatc aattcaagag ccatgtctgc atcataattg tgggagtgga ggaggctttt 1020
gttgctaggg aaggaatgcc ttcttagttc atggctttgg acttcggaca aggagcgcac 1080
agaatggggt taccattttt gaaaaaaatt acatttgaac cctccaacta ttatcatgta 1140
tgtttaatct acaatcctcg tccgctagaa gaagtttggg ttcaaagtca tccgctcaat 1200
gtaaacctg gagaggcagg gactaattga aaatagtatg ttagttggag ggtctgcatg 1260
tattatgtcc aaacattctc ttattattcc tgtatcatct ctgagaaatt catccgaaaa 1320
taataaaaca aaatggcctt ttttaaaaaa agaagctgat gcataggata ccaaagcgc 1380
cttgtccatt aggagcgtca gactttgaaa ataagaccaa taattccctg taagctatca 1440
tctcatcttt tattttgttt gaactttag acgtaggctt taagcgttcc atgatgttca 1500
gtcacatgtt gctgtctact tgattatgga atttaattca ttcggctcat aagaagataa 1560
aaggattatg acgttgaaga actctggtca ctcttactt acggtcacat aaaaacgatg 1620
catctttecc caccaaccat cttcaagtga acccatttc cttgcatga ggtaaggagt 1680
atgggttaag tcatcttcat gaaattagtc ccctagtgga gctaattcta ctactccat 1740
atttactcat tccactatat aacgccctca acgaccatcc tcaaagcaac ccaaacacct 1800
tcttctc 1807

<210> 5

52669.ST25

<211> 1494

<212> DNA

<213> Populus tremula x tremuloides

<220>

<221> misc_feature

<222> (337)..(337)

<223> unknown amino

<400> 5

tctgcttatg aatggcatgg acaacaagct ctgcattttc gccattttgc accacaaaat	60
cgttgtgtga agatttttga agggcaccag cacaactttg aaaagaactt gtgaaaagat	120
agctgggtcac cagatggaag caagggttact agtgcagatc gcatggctta tatatgaatc	180
tgtcaatgag tgtgtcttcc acctactgaa cctatctttg gatcatgcag cagcgacaaa	240
cagatttatac ttggggaaat ctgattacta ttttgtcagt ttgcttaaaa cgagttgtta	300
gtaacagtga ggatattygc gggcttgctt gtgagcnccc aacagaaaag attaaacagg	360
ttgtaatgga tactatgtta cattttactt aactttctca tatcctgctg acatgtttgc	420
agttcttttt tgcttgttgg cgatcctttt caactgaagt atgtacgcag gaatcaatta	480
gctcaaaatg tatgttcgag caatatgcct ggtgaatgtt tgtaagata cggaaaatta	540
agttactgaa cttgtactct ttgattttga aacaagtttc cgactatgtt cattgtccta	600
gcacgggtgat attgatacgt cgaattttca actatttttg tcttgttttc aactatttct	660
gtcttgattt tcaaagtctt tttgaaaata aaaaacccaa aaatatgttt tcagtgtctt	720
gttagctaaa tctgttgtag aaccaagcct ccccgacttg ctttgtgaaa aagaaagatg	780
ggagctgggg gttttgtggt gattatacaa aaccttacga gataactgat atacaagcaa	840
aaacaccctc atattattga atctgcgact tctacagttc atttctgagg aagaattagg	900
tgactacttt ctggactaga tatttccgtg ttcaatgcgt cgcaacaatt gatttggttg	960
ctggattatg aacatgataa tcacactata gaaagacgac gattatattt caagctatgg	1020
acaatgatgt tacctgctcg aagatcaata cgcttgaagg ccaagatatg atcascagg	1080
atgcaggcag tccagatggc ttcttgggat atgagcttca agattgttct tgaaaagaac	1140
caaataaatt gtaaaattct ctgggaatat tttttgttaa aattatcagc attgctattt	1200
tttttttcaa attcatttaa atattgctat tgggtagta gttcagcgca attaacttag	1260
ctggcagctt ggcattagct tcaaatcata tcacaacggt aaaagcagcc gttgcaacta	1320
gtcagcccac tagcatttgc tggtagccaa tcagatcata aaaggaaaac tagccgttgg	1380
caaaccctct ttcaacggtc atctccctgc ttcatgatat atatgcttag atatccag	1440
atccccctca acaaattatc tcaatcttca aaaccagtc cccagttaca gcag	1494

Sida 5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2004/000604

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C12N 15/82, C07K14/415

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2002138870 A1 (CHIANG ET AL), 26 Sept 2002 (26.09.2002), The whole document	1-7,9-17
A	--	8
A	WO 0070058 A2 (THE VICTORIA UNIVERSITY OF MANCHESTER), 23 November 2000 (23.11.2000), the whole document	1-17
A	Proc. Natl. Acad. Sci., Volume 95, October 1998, Fredrik Sterky et al, "Gene discovery in the wood-forming tissues of poplar: Analysis of 5,692 expressed sequence", pages 13330-13335, see the whole document	1-17
	--	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 July 2004

Date of mailing of the international search report

27-07-2004

Name and mailing address of the ISA/

Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

MALIN SÖDERMAN/BS

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2004/000604

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	The Plant Journal, Volume 16, No. 3, 1998, Magnus Hertzberg et al, "Molecular characterisation of a novel plant homeobox gene expressed in the maturing xylem zone of Populus tremula X tremuloides", pages 285-295, see whole document --	1-17
A	PNAS, Volume 98, No. 25, 2001, Magnus Hertzberg et al, "A transcriptional roadmap to wood formation", pages 14732-14737, see the whole document --	1-17
A	Proc. Natl. Acad. Sci., Volume 95, 1998, Wen-Jing Hu et al, "Compartmentalized expression of the two structurally and functionally distinct 4-coumarate: CoA ligase genes in aspen (Populus tremuloides), pages 5407-5412, see the whole document -- -----	1-17

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2004/000604

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of Item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
 - a. type of material
 - ☒ a sequence listing
 - ☐ table(s) related to the sequence listing
 - b. format of material
 - ☒ in written format
 - ☒ in computer readable form
 - c. time of filing/furnishing
 - ☐ contained in the international application as filed
 - ☒ filed together with the international application in computer readable form
 - ☐ furnished subsequently to this Authority for the purposes of search
2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2004/000604

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: **1-17**
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
see extra sheet
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2004/000604

Box II.2

The sequences on diskette (underlined sequences in figure 1a-1e) does not correspond to the sequences in claim 1 and figure 1a-1e (underlined sequences in figure 1a-1e and other sequence), thereby rendering the definition of the subject-matter of said claim unclear (Article 6 PCT).

Consequently, the search has been carried out for those parts of the application which appear to be clear, namely the sequences on diskette (underlined sequences in figure 1a-1e).

INTERNATIONAL SEARCH REPORT

Information on patent family members

30/04/2004

International application No.

PCT/SE 2004/000604

US	2002138870	A1	26/09/2002	US	2002124281	A	05/09/2002
				AU	8871601	A	22/03/2002
				AU	8876501	A	22/03/2002
				BR	0113654	A	23/09/2003
				CA	2421215	A	14/03/2002
				CN	1473198	T	04/02/2004
				EP	1320617	A	25/06/2003
				US	2002078474	A	20/06/2002
				WO	0220717	A	14/03/2002
				WO	0220812	A	14/03/2002

WO	0070058	A2	23/11/2000	AU	4934500	A	05/12/2000
				GB	9911379	D	00/00/0000
